

# A strategy for the computation of $m_b$ including $1/m$ terms



DESY 05-170  
SFB/CPP-05-58

**Michele Della Morte** \*

*Institut für Physik, Humboldt Universität, Newtonstr. 15, 12489 Berlin, Germany*  
E-mail: dellamor@physik.hu-berlin.de


**Nicolas Garron, Rainer Sommer**†

*DESY, Platanenallee 6, 15738 Zeuthen, Germany*  
E-mail: Nicolas.Garron@desy.de, Rainer.Sommer@desy.de

**Mauro Papinutto**

*NIC/DESY, Platanenallee 6, 15738 Zeuthen, Germany*  
E-mail: Mauro.Papinutto@desy.de

We consider HQET including the first order correction in  $1/m$ . A strategy for the computation of the b-quark mass following the scheme



is discussed. Only two quantities  $\Phi_{1/2}$  have to be considered in order to match QCD and HQET, since the spin-dependent interaction is easily eliminated due to the spin symmetry of the static theory. Quite simple formulae relate the renormalization group invariant b-quark mass ( $M_b$ ) to the B-meson mass. All entries in these formulae are non-perturbatively defined and can be computed in the continuum limit of the lattice regularized theory. For the numerically most critical part, we illustrate the cancellation of power divergences by a numerical example.

Numerical results for the  $1/m$  correction to  $M_b$ , are presented in a companion talk.

*XXIIIrd International Symposium on Lattice Field Theory*  
25-30 July 2005  
Trinity College, Dublin, Ireland

\*This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) in SFB Transregio 9 “Computational Particle Physics”.

†Speaker.

## 1. Introduction

Although HQET is the most natural effective theory for heavy-light systems, its lattice regularized version has practically only been used at lowest order. Indeed, a strategy to overcome the problem of power divergent mixings [1], was only found rather recently [2]. Its potential was demonstrated by a computation of the b-quark mass to lowest non-trivial order in  $1/m$ , the static approximation. Here we fill the formalism of [2], sketched in the abstract, with practicable definitions in terms of Schrödinger functional correlation functions and give a concrete formula for the  $1/m$ -correction to the quark mass.

Neglecting  $1/m^2$  corrections – as throughout this report – we write the HQET Lagrangian

$$\mathcal{L}_{\text{HQET}} = \mathcal{L}_{\text{stat}}(x) - \omega_{\text{spin}} \mathcal{O}_{\text{spin}}(x) - \omega_{\text{kin}} \mathcal{O}_{\text{kin}}(x) \quad (1.1)$$

$$\mathcal{O}_{\text{spin}} = \bar{\psi}_h \boldsymbol{\sigma} \mathbf{B} \psi_h, \quad \mathcal{O}_{\text{kin}} = \bar{\psi}_h \mathbf{D}^2 \psi_h, \quad (1.2)$$

such that the classical values for the coefficients are  $\omega_{\text{kin}} = \omega_{\text{spin}} = 1/(2m)$ . Since expectation values

$$\langle \mathcal{O} \rangle = \langle \mathcal{O} \rangle_{\text{stat}} + \omega_{\text{kin}} \langle \mathcal{O} \rangle_{\text{kin}} + \omega_{\text{spin}} \langle \mathcal{O} \rangle_{\text{spin}}, \quad (1.3)$$

$$\langle \mathcal{O} \rangle_{\text{kin}} = \sum_x \langle \mathcal{O} \mathcal{O}_{\text{kin}}(x) \rangle_{\text{stat}}, \quad \langle \mathcal{O} \rangle_{\text{spin}} = \sum_x \langle \mathcal{O} \mathcal{O}_{\text{spin}}(x) \rangle_{\text{stat}} \quad (1.4)$$

are defined through *insertions* of the higher dimensional terms  $\mathcal{O}_{\text{kin}}, \mathcal{O}_{\text{spin}}$  in the static theory, they are renormalizable by power counting. However, in order to have a well defined continuum limit the bare, dimensionful, couplings  $\omega_{\text{kin}}, \omega_{\text{spin}}$  have to be determined non-perturbatively [1, 2]. In the framework of lattice QCD, this is possible by matching a number of observables,  $\Phi_i$ ,  $i = 1 \dots n$ , between QCD and HQET, thus retaining the predictivity of QCD. It is essential to note that this matching can be carried out in a finite volume of linear extent  $L_1 \simeq 0.4 \text{ fm}$ , where heavy quarks can be simulated with a relativistic action [2, 3, 4].

Since the lowest order theory is spin-symmetric, it is trivial to form spin-averages which are independent of  $\omega_{\text{spin}}$ . One thus expects that  $n = 2$  is sufficient for a computation of the quark mass (in addition to  $\omega_{\text{kin}}$  there is an overall (state-independent) shift of energy levels, which we denote by  $m_{\text{bare}}$ ). For unexplained notation we refer to [2].

## 2. Basic observables

We consider the spin-symmetric combination

$$f_1^{\text{av}}(\theta, T) = Z_\zeta^4 \{f_1(\gamma_5)\}^{1/4} \{f_1(\gamma_1)\}^{3/4}, \quad (2.1)$$

formed from the boundary to boundary correlation functions

$$f_1(\Gamma) = -\frac{a^{12}}{2L^6} \sum_{\mathbf{u}, \mathbf{v}, \mathbf{y}, \mathbf{z}} \left\langle \bar{\zeta}_1'(\mathbf{u}) \Gamma \zeta_b'(\mathbf{v}) \bar{\zeta}_b(\mathbf{y}) \Gamma \zeta_l(\mathbf{z}) \right\rangle, \quad (2.2)$$

of the QCD Schrödinger functional of size  $T \times L^3$  and a periodicity phase  $\theta$  [5] for the quark fields. Replacing the b-quark field by the effective field  $\psi_h$ , using eq.(1.3,1.4), and accounting for the multiplicative renormalization of the boundary quark fields  $\zeta$ ,  $\bar{\zeta}$  one finds the  $1/m$  expansion

$$f_1^{\text{av}} = Z_{\zeta_h}^2 Z_\zeta^2 e^{-m_{\text{bare}} T} \{f_1^{\text{stat}} + \omega_{\text{kin}} f_1^{\text{kin}}\}, \quad (2.3)$$

where the aforementioned energy shift  $m_{\text{bare}}$  enters. Deviating from the choice in [2], we now define<sup>1</sup>

$$\Phi_1(L, M) = \ln(f_1^{\text{av}}(\theta, T)/f_1^{\text{av}}(\theta', T)) - \ln(f_1^{\text{stat}}(\theta, T)/f_1^{\text{stat}}(\theta', T)) \quad \text{for } T = L/2, \quad (2.5)$$

$$\Phi_2(L, M) = \frac{L}{2a} \ln(f_1^{\text{av}}(\theta, T-a)/f_1^{\text{av}}(\theta, T+a)), \quad (2.6)$$

with the expansion

$$\Phi_1(L, M) = \omega_{\text{kin}} R_1^{\text{kin}}, \quad \Phi_2(L, M) = L (m_{\text{bare}} + \Gamma_1^{\text{stat}} + \omega_{\text{kin}} \Gamma_1^{\text{kin}}) \quad (2.7)$$

$$R_1^{\text{kin}} = \frac{f_1^{\text{kin}}(\theta, T)}{f_1^{\text{stat}}(\theta, T)} - \frac{f_1^{\text{kin}}(\theta', T)}{f_1^{\text{stat}}(\theta', T)}, \quad (2.8)$$

$$\Gamma_1^{\text{stat}} = \frac{1}{2a} \ln(f_1^{\text{stat}}(\theta, T-a)/f_1^{\text{stat}}(\theta, T+a)), \quad (2.9)$$

$$\Gamma_1^{\text{kin}} = \frac{1}{2a} \left( \frac{f_1^{\text{kin}}(\theta, T-a)}{f_1^{\text{stat}}(\theta, T-a)} - \frac{f_1^{\text{kin}}(\theta, T+a)}{f_1^{\text{stat}}(\theta, T+a)} \right). \quad (2.10)$$

### 3. Step scaling functions

We choose  $L_1 \approx 0.4\text{fm}$ , where a computation of  $\Phi_i(L_1, M_b)$  is possible in lattice QCD (while at significantly larger values,  $L_1/a$  would have to be too large in order to control  $a^2$  effects). From eq. (2.7) one then gets  $\omega_{\text{kin}}, m_{\text{bare}}$  for lattice spacings  $a = \frac{a}{L_1} \times 0.4\text{fm}$ . On the other hand, contact to physical observables, e.g. the B-meson mass is made in large volume, where finite size effects are exponentially small. For reasonable values  $a/L_1 = 1/12$  and  $L_\infty \simeq 1.5\text{fm}$  at the same lattice spacing, one needs  $L_\infty/a \sim 50$ . This situation is avoided by first computing step scaling functions which connect  $\Phi_i(L_1, M)$  to  $\Phi_i(L_2, M), L_2 = 2L_1$  and then connecting to large volume.

With the Schrödinger functional coupling,  $u = \bar{g}^2(L)$ , everywhere, the continuum step scaling functions  $\sigma$  are defined by

$$\Phi_1(2L, M) = \sigma_1^{\text{kin}}(u) \Phi_1(L, M), \quad \sigma_1^{\text{kin}}(u) = \lim_{a/L \rightarrow 0} \frac{R_1^{\text{kin}}(2L)}{R_1^{\text{kin}}(L)} \Big|_{u=\bar{g}^2(L)} \quad (3.1)$$

and

$$\begin{aligned} \Phi_2(2L, M) - 2\Phi_2(L, M) &= \sigma_m(u) + [\omega_{\text{kin}} 2L (\Gamma_1^{\text{kin}}(2L) - \Gamma_1^{\text{kin}}(L))] \\ &= \sigma_m(u) + \sigma_2^{\text{kin}}(u) \Phi_1(L, M), \quad \sigma_2^{\text{kin}}(u) = \lim_{a/L \rightarrow 0} 2L \frac{\Gamma_1^{\text{kin}}(2L) - \Gamma_1^{\text{kin}}(L)}{R_1^{\text{kin}}(L)} \Big|_{u=\bar{g}^2(L)}. \end{aligned} \quad (3.2)$$

Here the static step scaling function

$$\sigma_m(u) = \lim_{a/L \rightarrow 0} 2L [\Gamma_1^{\text{stat}}(2L) - \Gamma_1^{\text{stat}}(L)]_{u=\bar{g}^2(L)}, \quad (3.3)$$

is not identical to  $\sigma_m(u)$  defined earlier [2], since  $\Gamma_1^{\text{stat}}$  differs from  $\Gamma^{\text{stat}}$  defined there. Note that the step scaling functions are independent of  $M$ , but  $\Phi_i(L, M)$  have a mass dependence from fixing  $\Phi_i(L_1, M)$  in the full theory.

<sup>1</sup>In the static computation of [2] the logarithmic derivative  $\Gamma$  of the correlation function  $f_A$  of the axial current with a boundary operator was used as a quantity to match effective theory and QCD. Including  $1/m$  terms its expansion reads

$$f_A = Z_A^{\text{HQET}} Z_\zeta Z_\zeta e^{-m_{\text{bare}} x_0} \left\{ f_A^{\text{stat}} + c_A^{\text{HQET}} f_{\delta A}^{\text{stat}} + \omega_{\text{kin}} f_A^{\text{kin}} + \omega_{\text{spin}} f_A^{\text{spin}} \right\}, \quad (2.4)$$

with the term  $f_{\delta A}^{\text{stat}}$  due to the  $1/m$  correction to the static axial current. While  $\omega_{\text{spin}}$  represents no problem, an extra observable is needed to fix  $f_{\delta A}^{\text{stat}}$ . Here, we avoid this complication by working exclusively with  $f_1^{\text{av}}$ .

#### 4. Large volume

The connection of  $\Phi_i$  to the spin-averaged B-meson mass,  $m_B$ , is

$$\begin{aligned} Lm_B - \Phi_2(L, M) &= [L(E^{\text{stat}} - \Gamma_1^{\text{stat}}(L))] + [L\omega_{\text{kin}}(\hat{E}^{\text{kin}} - \Gamma_1^{\text{kin}}(L))] \\ &= [L(E^{\text{stat}} - \Gamma_1^{\text{stat}}(L))] + \rho(u)\Phi_1(L, M), \quad \rho(u) = \lim_{a/L \rightarrow 0} L \frac{\hat{E}^{\text{kin}} - \Gamma_1^{\text{kin}}(L)}{R_1^{\text{kin}}(L)} \Big|_{u=\bar{g}^2(L)}. \end{aligned} \quad (4.1)$$

Here we have used the abbreviations

$$E^{\text{stat}} = \lim_{L \rightarrow \infty} \Gamma_1^{\text{stat}}(L), \quad \hat{E}^{\text{kin}} = \lim_{L \rightarrow \infty} \Gamma_1^{\text{kin}}(L), \quad (4.2)$$

where  $E^{\text{stat}}$  is the (unrenormalized) energy in large volume in the spin-averaged B-channel in static approximation and  $\omega_{\text{kin}} \hat{E}^{\text{kin}}$  is its  $1/m$  correction. The hat on  $\hat{E}^{\text{kin}}$  is to remind us that this quantity turns into an energy only upon multiplication with the dimensionful  $\omega_{\text{kin}}$ . Its numerical evaluation has already been investigated in [6]. We use [...] braces to indicate combinations which have a continuum limit by themselves. For example, the two terms in eq. (4.1) can be computed with different regularizations if this is useful.

#### 5. Final equation

The above equations are now easily combined to yield the  $1/m$  correction,  $m_B^{(1)}$ , to the (spin-averaged) B-meson mass via ( $L_2 = 2L_1$ ),

$$m_B = m_B^{\text{stat}} + m_B^{(1)} = m_B^{\text{stat}} + m_B^{(1a)} + m_B^{(1b)}, \quad (5.1)$$

$$L_2 m_B^{\text{stat}}(M) = [L_2(E^{\text{stat}} - \Gamma_1^{\text{stat}}(L_2))] + \sigma_m(u_1) + 2\Phi_2(L_1, M) \quad (5.2)$$

$$L_2 m_B^{(1a)}(M) = \sigma_2^{\text{kin}}(u_1)\Phi_1(L_1, M), \quad u_i = \bar{g}^2(L_i) \quad (5.3)$$

$$L_2 m_B^{(1b)}(M) = [L_2(\hat{E}^{\text{kin}} - \Gamma_1^{\text{kin}}(L_2))\omega_{\text{kin}}] = \rho(u_2)\sigma_1^{\text{kin}}(u_1)\Phi_1(L_1, M).$$

Again, terms in braces have a continuum limit. While  $m_B^{(1a)}$  is purely derived from finite volume, the term  $m_B^{(1b)}$  involves a large volume computation.

Starting from  $M_b^{\text{stat}}$ , the solution of the leading order equation,

$$m_B^{\text{exp}} = m_B^{\text{stat}}(M_b^{\text{stat}}), \quad (5.4)$$

and the slope

$$S = \frac{d}{dM} m_B^{\text{stat}} \Big|_{M=M_b^{\text{stat}}} = \frac{1}{L_1} \frac{d}{dM} \Phi_2(L_1, M) \Big|_{M=M_b^{\text{stat}}}, \quad (5.5)$$

we finally obtain the first order correction  $M_b^{(1)}$  to the RGI b-quark mass

$$M_b = M_b^{\text{stat}} + M_b^{(1)}, \quad M_b^{(1)} = -\frac{1}{S} m_B^{(1)}. \quad (5.6)$$

The final uncertainty for  $M_b$  due to the  $1/m$  expansion is of order  $O(\Lambda_{\text{QCD}}^3/M_b^2)$ , which translates into a numerical estimate of MeV scale. It is thus clear that other sources of error will dominate in a

practical calculation. Note that the precise value for  $m_B$  matters. One should use the spin-averaged mass  $m_B^{\text{experimental}} = \frac{1}{4}m_{B^0} + \frac{3}{4}m_{B^*} = [\frac{1}{4}5279 + \frac{3}{4}5325]\text{MeV} = 5314\text{MeV}$  if one can extrapolate  $E$  to the chiral limit of the light quark or

$$m_B^{\text{experimental}} = m_{B_s} + \frac{3}{4}m_{B_s^*} - \frac{3}{4}m_{B_0} = [5370 + \frac{3}{4}(5325 - 5279)]\text{MeV} = 5405\text{MeV} \quad (5.7)$$

if one works directly with a strange quark (as light quark). The latter formula neglects the dependence of the spin splitting on the light quark mass.

## 6. Remarks

The following facts are worth noting.

- The  $1/m$  expansion in heavy light systems is an expansion in terms of  $\Lambda_{\text{QCD}}/m$ , where all external scales have to be of order  $\Lambda_{\text{QCD}}$ . This applies in particular to our scale  $L_1^{-1}$ . Indeed, numerically it is rather close to  $\Lambda_{\text{QCD}}$  and explicit investigations [3, 4] have shown that the  $1/m$ -expansion is well behaved even when  $L^{-1}$  is a factor two larger.
- In our static computation [2, 3], we made the more natural choice  $\Gamma$  instead of  $\Gamma_1$ . Although it is advantageous to use  $\Gamma_1$  when one includes the  $1/m$  terms, the strategy can easily be formulated with  $\Gamma$ , at the expense of introducing a third quantity  $\Phi_i$  to fix  $c_A^{\text{HQET}}$ . Since this will certainly be required for the computation of the  $1/m$ -correction to  $F_B$ , we will follow also that approach.
- Note that at each order  $k$  in the expansion, the result is ambiguous by terms of order  $1/m^{k+1}$ . Thus both  $M_b^{(1)}$  and  $M_b^{\text{stat}}$  have an order  $1/m$  ambiguity (e.g. they change when  $L_1$  is changed), while in their sum  $M_b = M_b^{\text{stat}} + M_b^{(1)}$  the ambiguity is reduced to  $1/m^2$ .
- In the present formulation of the effective theory, the  $1/m$ -terms approach the continuum with an asymptotic rate  $\propto a$ , in contrast to the leading order terms where this is  $\propto a^2$  [2].
- Let us comment just on one numerical result at that point. The computation of  $\sigma_2^{\text{kin}}(u_1)$ , eq. (3.2), involves the difference of  $\Gamma_1^{\text{kin}}(2L) - \Gamma_1^{\text{kin}}(L)$ , where power divergent contributions cancel. As a typical case we choose  $L/a = 12$ ,  $T/a = 6$ , and the static action HYP2 (see [7]), where our simulations yield  $a^2\Gamma_1^{\text{kin}}(2L) = 0.5631(6)$ ,  $a^2\Gamma_1^{\text{kin}}(L) = 0.5595(2)$ , demonstrating a considerable cancellation. A detailed account of numerical results is presented in [8].

## References

- [1] L. Maiani, G. Martinelli and C.T. Sachrajda, Nucl. Phys. B368 (1992) 281,
- [2] ALPHA, J. Heitger and R. Sommer, JHEP 02 (2004) 022, hep-lat/0310035,
- [3] ALPHA, J. Heitger and J. Wennekers, JHEP 02 (2004) 064, hep-lat/0312016,
- [4] ALPHA, J. Heitger, A. Jüttner, R. Sommer and J. Wennekers, JHEP 11 (2004) 048, hep-ph/0407227,
- [5] S. Sint and R. Sommer, Nucl. Phys. B465 (1996) 71, hep-lat/9508012.
- [6] S. Dürr, A. Jüttner, J. Rolf and R. Sommer, (2004), hep-lat/0409058,
- [7] M. Della Morte, A. Shindler and R. Sommer, JHEP 08 (2005) 051, hep-lat/0506008,
- [8] ALPHA, M. Della Morte, N. Garron, M. Papinutto and R. Sommer, (2005), PoS (LAT2005) 224.